



Patient: ALBANA QEMALI DOB: November 13, 2013 Sex: F MRN: B000108750

# 3300 Organic Acids - Urine

63 Zillicoa Street Asheville, NC 28801 © Genova Diagnostics

## Order Number: P9140050

Reported: July 29, 2021 Received: July 14, 2021 Collected: July 12, 2021

# Organic Acids NUTRITIONAL

University Clinic of Gynecology and Obstetrics, Skopj Dimitar Kalcovski 50 Divizija 40/1-4 Skopje, 1000 Macedonia



Nutrient Need Overview						
Nutrient Need		Suggested	Provider			
0 1 2 3 4 5 6 7 8 9 10	DRI	Recommendations	Recommendations			
	0.6 mg	2 mg				
▲ ■ ■ ■ ■	0.6 mg	5 mg				
	8 mg	10 mg				
	0.6 mg	2 mg				
	12 mcg	50 mcg				
•	200 mcg	200 mcg				
<b>*</b>	1.2 mcg	10 mcg				
	130 mg	150 mg				
•	1.5 mg	1.5 mg				
	5 mg	5 mg				
<b>(</b>		5,000 IU				
•		25 billion CFU				
	Nutrient Need       Nutrient Need         0       1       2       3       4       5       6       7       8       9       10         •       <	Nutrient Need       DRI         0       1       2       3       4       5       6       7       8       9       10         0       1       2       3       4       5       6       7       8       9       10       DRI         0       1       2       3       4       5       6       7       8       9       10       DRI         0       1       2       3       4       5       6       7       8       9       10       DRI         0       0       1       0       6       mg       12       mg       12       mg       1.2       mg       1.5       mg       1.5       5       5       5       5       5       5       5       5       5       5       5       5       5       5       5       5       5       5       5	Nutrient Need         Suggested Recommendations           0         1         2         3         4         5         6         7         8         9         10         DRI         Suggested Recommendations           0         1         2         3         4         5         6         7         8         9         10         DRI         Suggested Recommendations           0         1         2         3         4         5         6         7         8         9         10         DRI         Suggested Recommendations           0         6         mg         0.6         mg         2         mg         5         mg         10         mg         2         mg         50         mcg         200         mcg         200         mcg         10         mcg         10         mg         10         mg         15         mg         1.5         mg         1.5         mg         5         mg <t< td=""></t<>			

Recommendations for age and gender-specific supplementation are set by comparing levels of nutrient functional need to optimal levels as described in the peer-reviewed literature. They are provided as guidance for short-term support of nutritional deficiencies only.

The Nutrient Need Overview is provided at the request of the ordering practitioner. Any application of it as a therapeutic intervention is to be determined by the ordering practitioner.

Page 2

# Page 3

# **Interpretation At-A-Glance**

# **Antioxidant Needs**

#### Glutathione



- Glutathione (GSH) is composed of cysteine, glutamine & glycine. GSH is a source of sulfate and plays a key role in antioxidant activity and detoxification of toxins.
- GSH requirement is increased with high-fat diets, cigarette smoke, cystinuria, chronic alcoholism, chronic acetaminophen use, infection, inflammation and toxic exposure.
- Deficiency may result in oxidative stress & damage, impaired detoxification, altered immunity, macular degeneration and increased risk of chronic illness.
- Food sources of GSH precursors include meats, poultry, fish, soy, corn, nuts, seeds, wheat germ, milk and cheese.



B1 is a required cofactor for enzymes involved in energy production from food,

Low B1 can result from chronic alcoholism, diuretics, digoxin, oral contraceptives and HRT, or large amounts of tea & coffee (contain anti-B1 factors).

B1 deficiency may lead to dry beriberi (e.g., neuropathy, muscle weakness),

wet beriberi (e.g., cardiac problems, edema), encephalopathy or dementia.

B2 is a key component of enzymes involved in antioxidant function, energy

Low B2 may result from chronic alcoholism, some anti-psychotic medications,

B2 deficiency may result in oxidative stress, mitochondrial dysfunction, low uric acid, low B3 or B6, high homocysteine, anemia or oral & throat inflammation.

Food sources include milk, cheese, eggs, whole grains, beef, chicken, wheat

B3 is used to form NAD and NADP, involved in energy production from food,

Low B3 may result from deficiencies of tryptophan (B3 precursor), B6, B2 or Fe

B3 deficiency may result in pellagra (dermatitis, diarrhea, dementia), neurologic

symptoms (e.g., depression, memory loss), bright red tongue or fatigue. Food sources include poultry, beef, organ meats, fish, whole grains, peanuts,

seeds, lentils, brewer's yeast and lima beans

(cofactors in B3 production), or from long-term isoniazid or oral contraceptive

production, detoxification, methionine metabolism and vitamin activation.

oral contraceptives, tricyclic antidepressants, quinacrine or adriamycin.

germ, fish, broccoli, asparagus, spinach, mushrooms and almonds.

fatty acid & cholesterol synthesis, cell signaling, DNA repair & cell

meats, brewer's yeast, blackstrap molasses, spinach, milk & eggs.

Food sources include lentils, whole grains, wheat germ, Brazil nuts, peas, organ

and for the synthesis of ATP, GTP, DNA, RNA and NADPH.

Thiamin - B1

**Riboflavin - B2** 

Niacin - B3

differentiation

use

Page 4

# Interpretation At-A-Glance

# B-Vitamin Needs

### Pyridoxine - B6



- B6 (as P5P) is a cofactor for enzymes involved in glycogenolysis & gluconeogenesis, and synthesis of neurotransmitters, heme, B3, RBCs and nucleic acids.
- Low B6 may result from chronic alcoholism, long-term diuretics, estrogens (oral contraceptives and HRT), anti-TB meds, penicillamine, L-DOPA or digoxin.
- B6 deficiency may result in neurologic symptoms (e.g., irritability, depression, seizures), oral inflammation, impaired immunity or increased homocysteine.
- Food sources include poultry, beef, beef liver, fish, whole grains, wheat germ, soybean, lentils, nuts & seeds, potato, spinach and carrots.

#### Biotin - B7



- Biotin is a cofactor for enzymes involved in functions such as fatty acid synthesis, mitochondrial FA oxidation, gluconeogenesis and DNA replication & transcription.
- Deficiency may result from certain inborn errors, chronic intake of raw egg whites, long-term TPN, anticonvulsants, high-dose B5, sulfa drugs & other antibiotics.
- Low levels may result in neurologic symptoms (e.g., paresthesias, depression), hair loss, scaly rash on face or genitals or impaired immunity.
- Food sources include yeast, whole grains, wheat germ, eggs, cheese, liver, meats, fish, wheat, nuts & seeds, avocado, raspberries, sweet potato and cauliflower.

#### Folate - B9

- Folate plays a key role in coenzymes involved in DNA and SAMe synthesis, methylation, nucleic acids & amino acid metabolism and RBC production.
- Low folate may result from alcoholism, high-dose NSAIDs, diabetic meds, H2 blockers, some diuretics and anti-convulsants, SSRIs, methotrexate, trimethoprim, pyrimethamine, triamterene, sulfasalazine or cholestyramine.
- Folate deficiency can result in anemia, fatigue, low methionine, increased homocysteine, impaired immunity, heart disease, birth defects and CA risk.
- Food sources include fortified grains, green vegetables, beans & legumes.

#### Cobalamin - B12



()

- B12 plays important roles in energy production from fats & proteins, methylation, synthesis of hemoglobin & RBCs, and maintenance of nerve cells. DNA & RNA.
- Low B12 may result from alcoholism, malabsorption, hypochlorhydria (e.g., from atrophic gastritis, H. pylori infection, pernicious anemia, H2 blockers, PPIs), vegan diets, diabetic meds, cholestyramine, chloramphenicol, neomycin or colchicine.
- B12 deficiency can lead to anemia, fatigue, neurologic symptoms (e.g., paresthesias, memory loss, depression, dementia), methylation defects or chromosome breaks.
- Food sources include shellfish, red meat, poultry, fish, eggs, milk and cheese.

#### KEY

Function of Nutrient

Cause of Deficiency

Complications of Deficiency

Food Sources of Nutrient

Magnesium



# **Interpretation At-A-Glance**

# **Mineral Needs**

4

#### Manganese

- Magnesium is involved in >300 metabolic reactions. Key areas include energy production, bone & ATP formation, muscle & nerve conduction and cell signaling.
- Deficiency may occur with malabsorption, alcoholism, hyperparathyroidism, renal disorders (wasting), diabetes, diuretics, digoxin or high doses of zinc.
- Low Mg may result in muscle weakness/spasm, constipation, depression, hypertension, arrhythmias, hypocalcemia, hypokalemia or personality changes.
- Food sources include dark leafy greens, oatmeal, buckwheat, unpolished grains, chocolate, milk, nuts & seeds, lima beans and molasses.

0

- Manganese plays an important role in antioxidant function, gluconeogenesis, the urea cycle, cartilage & bone formation, energy production and digestion.
- Impaired absorption of Mn may occur with excess intake of Fe, Ca, Cu, folic acid, or phosphorous compounds, or use of long-term TPN, Mg-containing antacids or laxatives.
- Deficiency may result in impaired bone/connective tissue growth, glucose & lipid dysregulation, infertility, oxidative stress, inflammation or hyperammonemia.
- Food sources include whole grains, legumes, dried fruits, nuts, dark green leafy vegetables, liver, kidney and tea.

# Zinc

- Zinc plays a vital role in immunity, protein metabolism, heme synthesis, growth & development, reproduction, digestion and antioxidant function.
- Low levels may occur with malabsorption, alcoholism, chronic diarrhea, diabetes, excess Cu or Fe, diuretics, ACE inhibitors, H2 blockers or digoxin.
- Deficiency can result in hair loss and skin rashes, also impairments in growth & healing, immunity, sexual function, taste & smell and digestion.
- Food sources include oysters, organ meats, soybean, wheat germ, seeds, nuts, red meat, chicken, herring, milk, yeast, leafy and root vegetables.





# Interpretation At-A-Glance

# **Microbiome & Digestive Support**

#### **Microbiome Support/Probiotics**



Probiotics have many functions. These include: production of some B vitamins and vitamin K; enhance digestion & absorption; decrease severity of diarrheal illness; modulate of immune function & intestinal permeability.

Alterations of gastrointestinal microflora may result from C-section delivery, antibiotic use, improved sanitation, decreased consumption of fermented foods and use of certain drugs.

 Some of the diseases associated with microflora imbalances include: IBS, IBD, fibromyalgia, chronic fatigue syndrome, obesity, atopic illness, colic and cancer.

Food sources rich in probiotics are yogurt, kefir and fermented foods.

#### **Digestive Support/Enzymes**



- Pancreatic enzymes are secreted by the exocrine glands of the pancreas and include protease/peptidase, lipase and amylase.
- Pancreatic exocrine insufficiency may be primary or secondary in nature. Any indication of insufficiency warrants further evaluation for underlying cause (i.e., celiac disease, small intestine villous atrophy, small bowel bacterial overgrowth).
- A high functional need for digestive enzymes suggests that there is an impairment related to digestive capacity.
- Determining the strength of the pancreatic enzyme support depends on the degree of functional impairment. Supplement potency is based on the lipase units present in both prescriptive and non-prescriptive agents.

# **Functional Imbalances**

### **Mitochondrial Dysfunction**



8

Mitochondria are a primary site of generation of reactive oxygen species. Oxidative damage is considered an important factor in decline of physiologic function that occurs with aging and stress.

Mitochondrial defects have been identified in cardiovascular disease, fatigue syndromes, neurologic disorders such as Parkinson's and Alzheimer's disease, as well as a variety of genetic conditions. Common nutritional deficiencies can impair mitochondrial efficiency.

#### **Toxic Exposure**

#### 

Methyl tert-Butyl Ether (MTBE) is a common gasoline additive used to increase octane ratings, and has been found to contaminate ground water supplies where gasoline is stored. Inhalation of MTBE may cause nose and throat irritation, as well as headaches, nausea, dizziness and mental confusion. Animal studies suggest that drinking MTBE may cause gastrointestinal irritation, liver and kidney damage and nervous system effects.

Styrene is classified by the US EPA as a "potential human carcinogen," and is found widely distributed in commercial products such as rubber, plastic, insulation, fiberglass, pipes, food containers and carpet backing.

Levels of these toxic substances should be examined within the context of the body's functional capacity for methylation and need for glutathione.

#### **Need for Methylation**



- Methylation is an enzymatic process that is critical for both synthesis and inactivation. DNA, estrogen and neurotransmitter metabolism are all dependent on appropriate methylation activity.
- B vitamins and other nutrients (methionine, magnesium, selenium) functionally support catechol-O-methyltransferase (COMT), the enzyme responsible for methylation.



Function of Nutrient

Cause of Deficiency

Complications of Deficiency





# **Oxidative Stress & Mitochondrial Dysfunction**



### Patient: ALBANA QEMALI

ID: P9140050

All biomarkers reported in mmol/mol creatinine unless otherwise noted.

Methodology: GCMS, LC/MS/MS, Alkaline Picrate, Colorimetric

#### **Organic Acids** Malabsorption & Dysbiosis Markers Vitamin Markers Reference Range Reference Malabsorption Markers Branched-Chain Catabolites (B1, B2, B3, ALA) Range 2.1 0.6 Indoleacetic Acid <= 4.2 α-Ketoadipic Acid <= 2.1 0.06 <dl Phenylacetic Acid α-Ketoisovaleric Acid <= 0.85 <= 0.15 0.22 Dysbiosis Markers α-Ketoisocaproic Acid <= 0.91 17.0 <dl α-Keto-β-Methylvaleric Dihydroxyphenylpropionic Acid (DHPPA) <= 23 <= 7.0 Acid 0.54 1.2 3-Hydroxyphenylacetic Glutaric Acid <= 0.92 <= 9.2 Acid 1.2 15 4-Hydroxyphenylacetic Isovalerylglycine <= 54 <= 37 Acid 0.07 Methylation Markers (Folate, B12) Benzoic Acid <= 0.10 115 0.2 Formiminoglutamic Acid **Hippuric Acid** <= 921 <= 2.7 (FIGlu) 0.8 Yeast / Fungal Dysbiosis Markers Methylmalonic Acid <= 2.2 48 **Biotin Markers D-Arabinitol** <= 51 8.9 8 Citramalic Acid <= 5.3 3-Hydroxypropionic Acid 6-23 <dl 6 Tartaric Acid <= 20 3-Hydroxyisovaleric Acid <= 38 Cellular Energy & Mitochondrial Markers Neurotransmitter Metabolites Reference Range Reference Fatty Acid Metabolism Kynurenine Markers (Vitamin B6) Range 2.0 3.4 <= 9.2 Adipic Acid <= 5.0 Kynurenic Acid 1.6 17.7 Quinolinic Acid Suberic Acid <= 4 2 <= 11.6 0.19 Kynurenic / Quinolinic Carbohydrate Metabolism >= 0.46 Ratio 0.27 8 Xanthurenic Acid <= 1.07 12-39 Pyruvic Acid 3.5 Catecholamine Markers 3.7-14.6 Lactic Acid <dl 9.8 a-Hydroxybutyric Acid <= 0.80 Homovanillic Acid 1.8-8.6 1.0 3.2 β-OH-Butyric Acid <= 3.4 Vanilmandelic Acid 1.5-5.0 13 0.06 3-Methyl-4-OHβ-OH-β-Methylglutaric <= 19 0.07-0.41 Acid phenylglycol **Energy Metabolism** Serotonin Markers 17.5 381 Citric Acid 62-648 5-OH-indoleacetic Acid 6.4-24.3 24 Reference Range Toxin & Detoxification Markers cis-Aconitic Acid 13-33 81 38 Isocitric Acid 38-97 Pyroglutamic Acid 22-64 ٠ 13 0.18 α-Ketoglutaric Acid 12-55 α-Ketophenylacetic Acid <= 0.50 (from Styrene) 5.4 5.2 Succinic Acid 0.8-10.4 a-Hydroxyisobutyric Acid <= 8.7 (from MTBE) 1.9 0.56 Malic Acid <= 2.7

© Genova Diagnostics · A. L. Peace-Brewer, PhD, D(ABMLI), Lab Director · CLIA Lic. #34D0655571 · Medicare Lic. #34-8475

Orotic Acid

Organic Acid Reference Ranges are Age Specific

0.38-0.91

Page 8

## Patient: ALBANA QEMALI

ID: P9140050

Page 9

Methodology: Colorimetric, thiobarbituric acid reactive substances (TBARS), Alkaline Picrate, Hexokinase/G-6-PDH, HPLC, GC/MS





F	'age	e 10	

# Commentary

For more information regarding Organic Acids clinical interpretation, please refer to the Organic Acids Support Guide at www.gdx.net/nutrevalguide.

# Lab Comments

The performance characteristics of all assays have been verified by Genova Diagnostics, Inc. Unless otherwise noted with •, the assay has not been cleared by the U.S. Food and Drug Administration.

The **Reference Range** is a statistical interval representing 95% or 2 Standard Deviations (2 S.D.) of the reference range population. One Standard Deviation (1 S.D.) is a statistical interval representing ~68% of the reference population. Values between 1 and 2 S.D. are not necessarily abnormal. Clinical Correlation is suggested.

